

Lesson 17

The area of a Surface of Revolution

Initializations

```
> restart;  
with(plots):
```

17.1 Area of a Surface of Revolution

If the curve $y=f(x)$, $a \leq x \leq b$ is rotated about the x axis, then the area of the resulting surface is given by

$$2\pi \int_a^b f(x) \sqrt{1 + (f'(x))^2} dx$$

The mathematical details will be provided in class.

Examples

Example 17.1.1

Find the area of the surface of revolution obtained by revolving the curve

$$y = \sin x, 0 \leq x \leq \frac{3}{4}\pi$$

about the x axis.

Solution

Use the formula provided above.

```
> f:=x->sin(x);
```

$$f := x \rightarrow \sin(x) \quad (2.1.1.1)$$

```
> e1:=2*Pi*Int(f(x)*sqrt(1+D(f)(x)^2), x=0..3*Pi/4);
```

$$e1 := 2\pi \int_0^{\frac{3}{4}\pi} \sin(x) \sqrt{1 + \cos(x)^2} dx \quad (2.1.1.2)$$


```
> surface_area:=simplify(value(e1));  
evalf(surface_area);
```

$$\text{surface_area} := -\frac{1}{2} \pi (-2\sqrt{2} - 2 \ln(1 + \sqrt{2}) - \sqrt{3} + 2 \ln(2) - 2 \ln(\sqrt{2} + \sqrt{2} \sqrt{3}))$$

$$12.00117140 \quad (2.1.1.3)$$

It is possible to visualize the resulting solid of revolution. The plotting routine makes use of a parametrization of the surface. You will learn the details in Calculus III. At this time you are not required to be able to do this.

```
> plot3d([x, sin(x)*cos(u), sin(x)*sin(u)], x=0..3*Pi/4, u=
0..2*Pi, style=patch, orientation=[-60, 70], axes=boxed,
labels=[x, y, z], scaling=constrained);
```



>

Example 17.1.2

Rotate the curve

$$x = \frac{(y^2 + 2)^{\frac{3}{2}}}{3}, 1 \leq y \leq 2$$

about the x axis and compute the area of the resulting surface of revolution.

Solution

This is an example where integration over y is appropriate. If dS denotes the surface area of a slice of the surface in a plane perpendicular to the x axis, then

$$dS = 2\pi r ds$$

where r denotes the radius of the slice and ds denotes the length of the piece of arc that generates the slice.

Therefore, dS can be written as

$$dS = 2\pi f(x) \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

but also as

$$dS = 2\pi y \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$

We now make use of the second representation.

```
> g:=y->1/3*(y^2+2)^(3/2);
```

$$g := y \rightarrow \frac{1}{3} (y^2 + 2)^{3/2} \quad (2.1.2.1)$$

```
> e1:=2*Pi*Int(y*sqrt(1+D(g)(y)^2), y=1..2);
```

$$e1 := 2\pi \int_1^2 y \sqrt{(1+y^2)^2} dy \quad (2.1.2.2)$$

```
> e2:=simplify(e1);
```

$$e2 := 2\pi \int_1^2 y (1+y^2) dy \quad (2.1.2.3)$$

```
> surface_area:=value(e2);
```

```
evalf(surface_area);
```

$$surface_area := \frac{21}{2} \pi$$

$$32.98672287 \quad (2.1.2.4)$$

```
>
```