

Lesson 2

Symbolic Computation

Initializations

```
> restart;
```

2.1 General Symbolic Computation

The real power of Maple lies in its symbolic capabilities. A computer algebra system allows the user to perform mathematical computations on the computer screen much like they used to be performed with pencil and paper. We now explore some of these symbolic features.

Examples

Example 2.1.1

Differentiation of Expressions.

Define the expression $e^{2x} \sin 7x$ and compute its first and second derivatives.

```
> expr:=exp(2*x)*sin(7*x);  
expr := e2x sin(7 x) (2.1.1.1)
```

```
> der1:=diff(expr, x);  
der1 := 2 e2x sin(7 x) + 7 e2x cos(7 x) (2.1.1.2)
```

```
> der2:=diff(expr, x$2);  
der2 := -45 e2x sin(7 x) + 28 e2x cos(7 x) (2.1.1.3)
```

```
>
```

Example 2.1.2

Equation Solving.

We consider three very familiar problems.

Problem 1.

The quadratic equation.

```
> eq1:=a*x^2+b*x+c=0;  
eq1 := a x2 + b x + c = 0 (2.1.2.1)
```

```
> sol1:=solve(eq1, x);  
sol1 := -\frac{1}{2} \frac{b - \sqrt{b^2 - 4ac}}{a}, -\frac{1}{2} \frac{b + \sqrt{b^2 - 4ac}}{a} (2.1.2.2)
```

```
>
```

Problem 2.

Simultaneous linear equations.

```
> eq2:={3*x-4*y=1, 5+x+7*y=-3};  
      eq2 := {3 x - 4 y = 1, 5 + x + 7 y = -3}      (2.1.2.3)
```

```
> sol2:=solve(eq2, {x, y});  
      sol2 := {y = -1, x = -1}      (2.1.2.4)
```

```
>
```

Problem 3.

Numeric Solutions.


Most real life equations cannot be solved in closed form, but require numerical routines instead. Maple's numeric equation solver is called **fsolve**. In this example we compute the points of intersection of the curves

$$y = \frac{x}{5} \quad \text{and} \quad y = \sin x$$

First we define the curves and make a sketch.

```
> c1:=x/5;  
   c2:=sin(x);  
  
      c1 :=  $\frac{1}{5} x$   
      c2 := sin(x)      (2.1.2.5)
```

```
> plot({c1, c2}, x=-10..10);
```



Clearly, (0, 0) is a point of intersection and there are points of intersection on the intervals (-4, -2) and (2, 4).

```
> x1:=fsolve(c1=c2, x=-4..-2);  
      x1 := -2.595739080      (2.1.2.6)
```

```
> x2:=fsolve(c1=c2, x=2..4);  
      x2 := 2.595739080      (2.1.2.7)
```

```
>
```

We use the **subs** command to obtain the corresponding points of intersection in (x, y) format..

```
> P1:=subs(x=x1, [x, c1]);  
      (2.1.2.8)
```

```
P1 := [-2.595739080, -0.5191478160]
```

(2.1.2.8)

```
> P2:=subs(x=x2, [x, c1]);
```

```
P2 := [2.595739080, 0.5191478160]
```

(2.1.2.9)

```
>
```